

## EXAMINING THE RELATIONSHIP BETWEEN SEVERITY OF THROMBOCYTOPENIA AND THE PATTERN OF MALARIA: A HOSPITAL-BASED STUDY

Javeria Rauf Saeed<sup>1</sup>

### How to cite this article

Saeed JR. Examining the Relationship between Severity of Thrombocytopenia and the Pattern of Malaria: A Hospital-Based Study. J Gandhara Med Dent Sci. 2024;11(1):34-38

**Date of Submission:** 19-09-2023

**Date Revised:** 18-11-2023

**Date Acceptance:** 25-11-2023

### Correspondence

<sup>1</sup>Javeria Rauf Saeed, Assistant

Professor, Haematology, Saidu

Medical College, Swat

☎: +92-333-5031246

✉: [safwansyed258@gmail.com](mailto:safwansyed258@gmail.com)

<https://doi.org/10.37762/jgm.11-1.557>

### ABSTRACT

#### OBJECTIVES

To determine the association between the severity of thrombocytopenia and the pattern of malaria.

#### METHODOLOGY

This descriptive cross-sectional study included 194 patients who tested positive for malarial parasites on peripheral smear examination. The study comprised smear-positive malaria cases, both males and females, who were Pakistani nationals aged between 8 and 50. Exclusion criteria were individuals with dengue, typhoid, immunocompromised status, liver disease, and pregnancy. Data were recorded for age, gender, the pattern of malaria (*Plasmodium* (*P.*) *vivax*, *P. falciparum*, or mixed), and type of thrombocytopenia (mild, moderate, and severe). The association between the severity of thrombocytopenia and the pattern of malaria was determined using a chi-square test, with the analysis stratified by gender.

#### RESULTS

The mean age of the participants was 28.58±11.70 years, with 114 (58.76%) being male. The lowest platelet count (64,627 ± 43,062) was found in cases of falciparum malaria, while the highest mean platelet count was found in cases of vivax malaria (130,919 ± 107,723), and this difference was statistically significant ( $p=0.004$ ). The most common occurrence of severe thrombocytopenia was in falciparum malaria cases ( $n=11$ , 42.31%), and the difference was statistically significant ( $p=0.024$ ).

#### CONCLUSION

Falciparum malaria has the lowest mean platelet count and more severe thrombocytopenia than vivax and mixed type.

**KEYWORDS:** Falciparum Malaria, Malaria, Platelet Count, Thrombocytopenia, Plasmodium Vivax

## INTRODUCTION

Malaria ranks as the fifth major cause of death globally.<sup>1</sup> Malaria poses a significant public health threat to individuals living in Asian countries, leading to substantial illness and economic consequences.<sup>2</sup> Despite numerous control measures, around 40% of the population remains vulnerable to this infection. The primary cause of malaria is *Plasmodium* species, namely vivax, falciparum, malaria, and ovale.<sup>3</sup> Despite being categorized as a country with moderate malaria prevalence, Pakistan still has an alarming number of around 177 million people who remain at risk of being infected with the disease. Around 60% of Pakistan's population reside in regions where malaria is endemic.<sup>4</sup> Out of the total malaria burden in Pakistan, *Plasmodium vivax* accounts for 81.3%, *P. falciparum* for 14.7%, and mixed-species for 4%.<sup>4</sup> Malaria is linked to haematological complications like anaemia and thrombocytopenia, caused by factors such as haemolysis, parasitic infections, antimalarial drugs, and deficiencies in essential vitamins and minerals like iron,

folate, and vitamin B12.<sup>5,6</sup> Despite widespread attention from the scientific community regarding anaemia associated with malaria and its associated mortality, thrombocytopenia has received less attention. However, the presence of thrombocytopenia can serve as a helpful indicator for primary care physicians in clinical diagnosis when other tests may not be accessible.<sup>7,8</sup> While the exact mechanism responsible for thrombocytopenia in malaria is not yet fully comprehended, some researchers propose that the malarial parasite reduces the production of platelets from megakaryocytes as a means of survival, thereby evading platelet-mediated clearance.<sup>9</sup> A previous study found that 80% of malaria cases had thrombocytopenia, which was observed in varying degrees of severity: 36.25% with mild thrombocytopenia, 40.0% with moderate thrombocytopenia, and 23.75% with severe thrombocytopenia.<sup>10</sup> Understanding haematological parameters is essential for assessing the severity of malaria and avoiding invasive investigations. Thrombocytopenia is a common complication of malaria, and its severity has been observed to vary

across affected individuals. Exploring the relationship between the severity of thrombocytopenia and the pattern of malaria could provide valuable insights into the pathophysiology of this condition. Specifically, understanding the role of malaria contributes to the degree of thrombocytopenia in malaria may aid in the early identification and treatment of severe cases, thereby reducing morbidity and mortality rates associated with this condition. Studying this relationship could also inform the development of new therapeutic approaches for malaria that specifically target the mechanisms underlying thrombocytopenia in the disease. This study aimed to determine the association between the severity of thrombocytopenia and the pattern of malaria.

## METHODOLOGY

This descriptive cross-sectional study was conducted at the Department of Pathology, Saidu Group of Teaching Hospitals, Swat, from June 10, 2022, to February 28, 2023. Ethical approval was obtained from the hospital's ethical committee. Ethical approval was not required separately, as it had already been secured through the informed consent process. All patients had consented to using their records for research purposes. A total of 194 patients who tested positive for malarial parasite on peripheral smear examination were included in the analysis. The sample size was calculated using the WHO sample size calculator at a 95% confidence level, with a 6% margin of error, and assuming a frequency of severe thrombocytopenia of 23.75% among malaria-positive patients.<sup>10</sup> The study included smear-positive malaria cases, both male and female, Pakistani nationals between the ages of 8 and 50. Exclusion criteria consisted of individuals with dengue, typhoid, immunocompromised status, liver disease, and pregnancy (in all these cases, platelet count can be affected). Cases recruited in the study underwent a detailed history-taking and complete general and systemic examination using a proforma. A medical specialist examined all patients, followed by routine and specific blood investigations to evaluate falciparum/vivax malaria. The platelet counts were obtained using a fully automated quantitative analyzer (Mandfredi, Italy 820-I) and were categorized into three subgroups - mild thrombocytopenia, moderate thrombocytopenia, and severe thrombocytopenia - based on the platelet count reference of previous study.<sup>10</sup> Mild thrombocytopenia was characterized by platelet counts ranging from 100 to 150  $\times 10^3/\mu\text{l}$ , moderate thrombocytopenia from 50 to 100  $\times 10^3/\mu\text{l}$ , and severe thrombocytopenia by counts below 50  $\times 10^3/\mu\text{l}$ . For patients with *P. falciparum*, *P. vivax* malaria, and mixed infections, treatment was

administered according to the standard protocol, which varied depending on the clinical severity. The data analysis was performed using R programming version 4.1.2. Mean and standard deviation were computed for numerical data such as age and platelet count, while frequency and percentages were calculated for qualitative data such as gender, malaria pattern, and thrombocytopenia severity. The platelet count was compared among malaria types using an ANOVA test. The association between the severity of thrombocytopenia and the pattern of malaria was determined using a chi-square test, with the analysis stratified by gender. The level of significance was  $p < 0.05$ .


## RESULTS

The mean age of the participant was  $28.58 \pm 11.70$  years. Of total 194 participants, 114 (58.76%) were males and 80 (41.24%) were females. The participants were divided into three age groups: 8-20 ( $n=55$ , 28.35%), 21-35 ( $n=81$ , 41.75%), and 35-50 ( $n=58$ , 29.90%). Thrombocytopenia was absent in 39 (20.10%). Most common type of thrombocytopenia was moderate, found in 63 (32.47%), followed by mild ( $n=47$ , 24.23%) and the least was severe ( $n=45$ , 23.20%). Among the participants, 159 (81.96%) had vivax malaria, 26 (13.40%) had falciparum malaria, and 9 (4.64%) had mixed malaria. (Table 1). Fig 1 provides information on the mean platelet count and standard error for three categories of thrombocytopenia: mild, moderate, and severe. The mean platelet count ( $\times 10^3/\mu\text{l}$ ) for mild thrombocytopenia was 12.44 with a standard deviation (SD) of 2.256. The mean platelet count for moderate thrombocytopenia was 12.4471, with an SD of 1.864. Finally, the mean platelet count for severe thrombocytopenia was 26.707, with an SD of 10.915. The lowest mean platelet count ( $\times 10^3/\mu\text{l}$ ) was found for falciparum malaria, with a value of  $64.627 \pm 43.062$ , while the highest mean platelet count was found for vivax malaria, with a value of  $130.919 \pm 107.723$ . The mean platelet count ( $\times 10^3/\mu\text{l}$ ) for mixed malaria was  $79.143 \pm 39.132$ . The ANOVA test showed a significant difference in mean platelet count ( $\times 10^3/\mu\text{l}$ ) between the three types of malaria ( $p=0.004$ ). Further analysis using multiple comparison tests revealed that the only statistically significant difference was found between falciparum and mixed type ( $p=0.002$ ). (Table 2) Figure 2 shows the platelet count in different malaria patterns in both genders. The difference was statistically significant only in males. The data analysis revealed that the most common severe thrombocytopenia was found in falciparum malaria, with 11 cases representing 42.31% of the overall sample. The second most common was in mixed

malaria, with 3 cases representing 33.33%, and the least was found in vivax malaria (n=31, 19.5%). In contrast, moderate thrombocytopenia was most frequent in mixed malaria, with (n=4, 44.44%). The differences in severe and moderate thrombocytopenia incidence were statistically significant (p=0.024). Furthermore, the analysis showed significant differences between males (p=0.041) and females (p=0.034) in the incidence of severe and moderate thrombocytopenia with the type of malaria. (Table 3)

**Table 1: Distribution of Gender, Ages, Thrombocytopenia and Malarial Pattern**


Variable	Characteristic	N = 194 <sup>1</sup>
Gender	Female	80 (41.24)
	Male	114 (58.76)
Age Group (years)	8-20	55 (28.35)
	21-35	81 (41.75)
	35-50	58 (29.90)
Thrombocytopenia	Nil	39 (20.10)
	Mild	47 (24.23)
	Moderate	63 (32.47)
	Severe	45 (23.20)
Malaria	Falciparum	26 (13.40)
	Mixed	09 (4.64)
	Vivax	159 (81.96)



**Figure 1: Mean Platelet Count in Various Types of Thrombocytopenia**

**Table 2: Comparison of Platelet Count Among Various Patterns of Malaria**

Characteristics	Falciparum (F), N=26 <sup>1</sup>	Mixed (M), N=9 <sup>1</sup>	Vivax (V), N=159 <sup>1</sup>	over all P-Value*	P-Values for Multiple Comparisons**
Platelet Count(x1000/µl)	64.627 ± 43.062	79.143 ± 39.132	130.919 ± 107.723	0.004	F vs M = 0.7 F vs V=0.002 M vs V=0.13



**Figure 2: Platelet Count in Different Patterns of Malaria in Both Genders**

**Table 3: Association of Severity of Thrombocytopenia and Pattern of Malaria**

Variable	Thrombocytopenia	Falciparum, N=26	Mixed, N=9	Vivax, N=159	P-Value*
Overall	Mild	06 (23.08)	02 (22.22)	39 (24.53)	0.024
	Moderate	09 (34.62)	04 (44.44)	50 (31.45)	
	Nil	0 (0.00)	0 (0.00)	39 (24.53)	
	severe	11 (42.31)	03 (33.33)	31 (19.50)	
Male	mild	05 (29.41)	01 (50.00)	22 (23.16)	0.041
	moderate	06 (35.29)	01 (50.00)	29 (30.53)	
	nil	0 (0.00)	0 (0.00)	22 (23.16)	
	severe	06 (35.29)	0 (0.00)	22 (23.16)	
Female	mild	01 (11.11)	01 (14.29)	17 (26.56)	0.034
	moderate	03 (33.33)	03 (42.86)	21 (32.81)	
	nil	0 (0.00)	0 (0.00)	17 (26.56)	
	severe	05 (55.56)	03 (42.86)	09 (14.06)	

\*chi-square test

**DISCUSSION**

This study aimed to determine the association between the severity of thrombocytopenia and the pattern of malaria. Our findings showed that falciparum malaria had the lowest mean platelet count, while vivax malaria had the highest. There was a significant difference in mean platelet count between the three types of malaria, with only falciparum and mixed type showing a statistically significant difference. The Falciparum malaria has more severe thrombocytopenia than vivax and mixed-type malaria. The mechanism underlying the observed differences in mean platelet count among the three types of malaria is not fully understood.<sup>11</sup> However, it is thought that falciparum malaria may induce more severe thrombocytopenia compared to vivax and mixed-type malaria due to various factors.<sup>12</sup> These include the destruction of platelets by infected

red blood cells, immune-mediated clearance of platelets, and the production of platelet antibodies. Additionally, falciparum malaria is known to induce endothelial dysfunction, which may also contribute to platelet destruction and thrombocytopenia.<sup>13</sup> Thrombocytopenia is common in individuals with acute malaria, irrespective of whether they are infected with *P. falciparum* or *P. vivax*. Although thrombocytopenia may also be a characteristic of viral febrile illnesses, in areas where malaria is prevalent, its presence is considered a diagnostic indicator.<sup>14</sup> Our study found that thrombocytopenia was present in 75% of patients diagnosed with malaria. This prevalence rate is consistent with previous studies conducted in India, which reported a range of 72.0% to 80.0%.<sup>10,15,16</sup> Specifically, Jadhav et al. reported a prevalence of 78.4%, while Shetty et al. found a prevalence of 72.0%.<sup>16</sup> These results suggest that thrombocytopenia is a common finding among patients with malaria in Asian countries and highlight the need for continued research to understand better the mechanisms underlying this phenomenon.<sup>15</sup> Thrombocytopenia is a common complication of malaria. Researchers have conducted extensive studies to understand the relationship between thrombocytopenia and malaria.<sup>10,15,16</sup> Our findings suggest that mild thrombocytopenia is more likely to occur in individuals with *P. vivax* malaria. In contrast, severe thrombocytopenia is more commonly linked to *P. falciparum* malaria, which is considered more severe than *P. vivax* malaria. Moreover, the severity of thrombocytopenia is significantly correlated with the presence of the M antigen. This antigen is a protein on the surface of malaria-infected red blood cells and is involved in binding infected cells to the blood vessel walls. Studies have shown that the presence of the M antigen is associated with the development of severe thrombocytopenia.<sup>10</sup> Other studies, including Memon et al., have reported similar outcomes, showing that hospitalized malaria patients with *P. falciparum* are more likely to develop severe thrombocytopenia.<sup>17</sup> However, Jadhav et al. reported that severe thrombocytopenia can occur in either type of malaria, and the presence of thrombocytopenia does not distinguish between them.<sup>16</sup> In contrast, Kaur et al. found that patients with *P. vivax* malaria can also experience severe thrombocytopenia, contradicting the previous notion that severe thrombocytopenia is primarily associated with *P. falciparum* malaria.<sup>18</sup> These findings suggest that the relationship between thrombocytopenia and malaria is complex and may depend on multiple factors, including the type of malaria, the severity of the infection, and the presence of specific antigens.

## LIMITATIONS

This study has limitations that need to be considered when interpreting the findings. The descriptive and cross-sectional study cannot establish a temporal relationship between thrombocytopenia and malaria. There were fewer cases of *P. falciparum* malaria included in the analysis than *P. vivax* malaria cases, which could limit the comparability and generalizability of the results. Other factors, such as sex, genetics, and immune status, may play a role in thrombocytopenia development, were not examined. The study may have also been affected by selection bias as it may have only included individuals with more severe malaria cases, skewing the findings. Further research is needed to confirm the relationship between thrombocytopenia and malaria and address these limitations.

## CONCLUSIONS

Our study suggests a significant difference in mean platelet count between the three types of malaria, with falciparum malaria having the lowest mean platelet count and more severe thrombocytopenia compared to vivax and mixed-type malaria. These findings highlight the importance of monitoring platelet count in individuals with malaria, especially those with falciparum malaria, as severe thrombocytopenia can lead to serious complications.

**CONFLICT OF INTEREST:** None

**FUNDING SOURCES:** None

## REFERENCES

1. Deribew A, Tessema GA, Deribe K, Melaku YA, Lakew Y, Amare AT, et al. Trends, causes, and risk factors of mortality among children under 5 in Ethiopia, 1990-2013: findings from the Global Burden of Disease Study 2013. *Popul Health Metr.* 2016;14(1):42.
2. Shah S, Abbas G, Riaz N, ur Rehman A, Hanif M, Rasool MF. Burden of communicable diseases and cost of illness: Asia pacific region. *Expert Rev Pharmacoecon Outcomes Res.* 2020;20(4):343-54.
3. Mohammed OIA, Abdalla MHA. Determination of Prothrombin Time, Activated Partial Thromboplastin Time and D-Dimer Levels among Malaria Infected Patients in Sinnar State, Sudan. *OALib.* 2019;06(12):1-6.
4. Qureshi NA, Fatima H, Afzal M, Khattak AA, Nawaz MA. Occurrence and seasonal variation of human Plasmodium infection in Punjab Province, Pakistan. *BMC Infect Dis.* 2019;19(1):935.
5. Awoke N, Arota A. Profiles of hematological parameters in Plasmodium falciparum and Plasmodium vivax malaria patients attending Tercha General Hospital, Dawuro Zone, South Ethiopia. *Infect Drug Resist.* 2019;12:521-7.
6. Latif I, Jamal A. Hematological changes in complete blood

- picture in paediatric patients of malaria caused by Plasmodium vivax and falciparum. J Ayub Med Coll Abbottabad. 2015;27(2):351-5. PMID 26411114.
7. Bahati YL, Delanghe J, Balaluka GB, Philippé J. Exploration of the relationship between anemia and iron and zinc deficiencies in children under 5 years of age living in the malaria endemic area of South Kivu/Democratic Republic of Congo. Ann Hematol. 2022;101(6):1181-9.
  8. Ibeji JU, Mwambi H, Iddrisu A-K. Bayesian spatiotemporal modelling and mapping of malaria and anaemia among children between 0 and 59 months in Nigeria. Malar J. 2022;21(1):1-12.
  9. Srivastava K, Sharma M, Mitchell WB. Malaria and thrombopoiesis: a possible mechanism for the malarial thrombocytopenia. J Immunol Infect Inflamm Dis. 2017;6:2.
  10. Kumar M, Kumar A, Panwar P, Kant R. Correlation of Presence and Severity of thrombocytopenia with Types and Severity of Malaria: A study from tertiary care center of North India. J Family Med Prim Care. 2022;11(7):3929-33.
  11. de Mast Qd, Groot E, Lenting PJ, de Groot PG, McCall M, Sauerwein RW, et al. Thrombocytopenia and release of activated von Willebrand Factor during early Plasmodium falciparum malaria. J Infect Dis. 2007;196(4):622-8.
  12. Araujo CF, Lacerda MV, Abdalla DS, Lima ES. The role of platelet and plasma markers of antioxidant status and oxidative stress in thrombocytopenia among patients with vivax malaria. Mem Inst Oswaldo Cruz. 2008;103(6):517-21.
  13. Gramaglia I, Sahlin H, Nolan JP, Frangos JA, Intaglietta M, van der Heyde HC. Cell- rather than antibody-mediated immunity leads to the development of profound thrombocytopenia during experimental Plasmodium berghei malaria. J Immunol. 2005;175(11):7699-707.
  14. Saberi N, Raeisi A, Gorouhi MA, Vatandoost H, Mashayekhi M, Nikpour F, et al. Current Situation of Malaria and Resistance of Main Vectors to WHO Recommended Insecticides in an Endemic Area, Southeastern Iran. J Med Entomol. 2022;59(2):607-14.
  15. Shetty G, Avabratha KS, Gonsalves S, Dany A, Rai BS. Thrombocytopenia in children with malaria—A study from coastal Karnataka, India. Asian Pac J Trop Dis. 2012;2(2):107-9.
  16. Jadhav UM, Patkar VS, Kadam NN. Thrombocytopenia in malaria--correlation with type and severity of malaria. J Assoc Physicians India. 2004;52:615-8. PMID 15847353.
  17. Memon AR, Afsar S. Thrombocytopenia in hospitalized malaria patients. Pak J Med Sci. 2006;22:141-3.
  18. Kaur D, Wasir V, Gulati S, Bagga A. Unusual presentation of Plasmodium vivax Malaria with severe thrombocytopenia and acute renal failure. J Trop Pediatr. 2007;53(3):210-2.

## CONTRIBUTORS

1. **Javeria Rauf Saeed** – *Concept & Design; Data Acquisition; Data Analysis/Interpretation; Drafting Manuscript; Critical Revision; Supervision; Final Approval*



LICENSE: JGMDS publishes its articles under a Creative Commons Attribution Non-Commercial Share-Alike license (CC-BY-NC-SA 4.0).  
 COPYRIGHTS: Authors retain the rights without any restrictions to freely download, print, share and disseminate the article for any lawful purpose.  
 It includes scholarly networks such as Research Gate, Google Scholar, LinkedIn, Academia.edu, Twitter, and other academic or professional networking sites.